

## **Blue diode laser versus traditional infrared diode laser and quantic molecular resonance scalpel: clinical and histological findings after excisional biopsy of benign oral lesions (Erratum)**

Margherita Gobbo  
Rossana Bussani  
Giuseppe Perinetti  
Katia Rupel  
Lorenzo Bevilacqua  
Giulia Ottaviani  
Matteo Biasotto

# Blue diode laser versus traditional infrared diode laser and quantic molecular resonance scalpel: clinical and histological findings after excisional biopsy of benign oral lesions (Erratum)

**Margherita Gobbo,<sup>a,b</sup> Rossana Bussani,<sup>b,c</sup> Giuseppe Perinetti,<sup>b</sup> Katia Rupel,<sup>a,b</sup> Lorenzo Bevilacqua,<sup>b</sup> Giulia Ottaviani,<sup>a,b,\*</sup> and Matteo Biasotto<sup>a,b</sup>**

<sup>a</sup>Division of Oral Medicine and Pathology, Dental Clinic, Ospedale Maggiore, Trieste, Italy

<sup>b</sup>University of Trieste, Department of Surgical, Medical and Health Sciences, Trieste, Italy

<sup>c</sup>Institute of Pathologic Anatomy, Ospedale di Cattinara, Trieste, Italy

[DOI: [10.1117/1.JBO.24.2.029803](https://doi.org/10.1117/1.JBO.24.2.029803)]

This article [*J. Biomed. Opt.* **22**(12), 121602 (2017), doi: [10.1117/1.JBO.22.12.121602](https://doi.org/10.1117/1.JBO.22.12.121602)] was originally published online on 12 July 2017 with an error in the spelling of an author's name, Lorenzo Bevilacqua.

This article was corrected online on 20 February 2019.