

PROCEEDINGS OF SPIE

***Harnessing Relativistic Plasma
Waves as Novel Radiation
Sources from Terahertz to
X-Rays and Beyond***

**Dino A. Jaroszynski
Antoine Rouse**
Editors

**21–23 April 2009
Prague, Czech Republic**

Sponsored by
SPIE Europe

Cooperating Organizations
Institute of Physics, Academy of Sciences (Czech Republic)
Department of X-Ray Lasers, Institute of Physics (Czech Republic)
Czech and Slovak Society for Photonics
Photonics Society of Poland (Poland)

Published by
SPIE

Volume 7359

Proceedings of SPIE, 0277-786X, v. 7359

SPIE is an international society advancing an interdisciplinary approach to the science and application of light.

The papers included in this volume were part of the technical conference cited on the cover and title page. Papers were selected and subject to review by the editors and conference program committee. Some conference presentations may not be available for publication. The papers published in these proceedings reflect the work and thoughts of the authors and are published herein as submitted. The publisher is not responsible for the validity of the information or for any outcomes resulting from reliance thereon.

Please use the following format to cite material from this book:

Author(s), "Title of Paper," in *Harnessing Relativistic Plasma Waves as Novel Radiation Sources from Terahertz to X-Rays and Beyond*, edited by Dino A. Jaroszynski, Antoine Rousse, Proceedings of SPIE Vol. 7359 (SPIE, Bellingham, WA, 2009) Article CID Number.

ISSN 0277-786X
ISBN 9780819476333

Published by

SPIE

P.O. Box 10, Bellingham, Washington 98227-0010 USA
Telephone +1 360 676 3290 (Pacific Time) · Fax +1 360 647 1445
SPIE.org

Copyright © 2009, Society of Photo-Optical Instrumentation Engineers

Copying of material in this book for internal or personal use, or for the internal or personal use of specific clients, beyond the fair use provisions granted by the U.S. Copyright Law is authorized by SPIE subject to payment of copying fees. The Transactional Reporting Service base fee for this volume is \$18.00 per article (or portion thereof), which should be paid directly to the Copyright Clearance Center (CCC), 222 Rosewood Drive, Danvers, MA 01923. Payment may also be made electronically through CCC Online at copyright.com. Other copying for republication, resale, advertising or promotion, or any form of systematic or multiple reproduction of any material in this book is prohibited except with permission in writing from the publisher. The CCC fee code is 0277-786X/09/\$18.00.

Printed in the United States of America.

Publication of record for individual papers is online in the SPIE Digital Library.

SPIE 
Digital Library

SPIDigitalLibrary.org

Paper Numbering: Proceedings of SPIE follow an e-First publication model, with papers published first online and then in print and on CD-ROM. Papers are published as they are submitted and meet publication criteria. A unique, consistent, permanent citation identifier (CID) number is assigned to each article at the time of the first publication. Utilization of CIDs allows articles to be fully citable as soon they are published online, and connects the same identifier to all online, print, and electronic versions of the publication. SPIE uses a six-digit CID article numbering system in which:

- The first four digits correspond to the SPIE volume number.
- The last two digits indicate publication order within the volume using a Base 36 numbering system employing both numerals and letters. These two-number sets start with 00, 01, 02, 03, 04, 05, 06, 07, 08, 09, 0A, 0B ... 0Z, followed by 10-1Z, 20-2Z, etc.

The CID number appears on each page of the manuscript. The complete citation is used on the first page, and an abbreviated version on subsequent pages. Numbers in the index correspond to the last two digits of the six-digit CID number.

Contents

- ix *Conference Committee*
- xi *Photon physics: from wave mechanics to quantum electrodynamics (Plenary Paper)*
[7355-100]
O. Keller, Aalborg Univ. (Denmark)

SESSION 1 TOWARDS A COMPACT X-FEL I

- 7359 02 **Towards a compact 0.1-10 MeV broadband betatron photon source (Invited Paper)**
[7359-01]
C. E. Clayton, Univ. of California, Los Angeles (United States); S. F. Martins, J. L. Martins, Instituto Superior Técnico (Portugal); D. K. Johnson, S. Wang, K. A. Marsh, Univ. of California, Los Angeles (United States); P. Muggli, M. J. Hogan, D. Walz, Univ. of Southern California (United States); R. A. Fonseca, Instituto Superior Técnico (Portugal); E. Oz, C. D. Barnes, C. L. O'Connell, I. Blumenfeld, N. Kirby, R. Ischebeck, Univ. of Southern California (United States); C. Huang, M. Zhou, W. Lu, Univ. of California, Los Angeles (United States); S. Deng, T. Katsouleas, Univ. of Southern California (United States); W. B. Mori, Univ. of California, Los Angeles (United States); R. H. Siemann, Univ. of Southern California (United States); L. O. Silva, Instituto Superior Técnico (Portugal); C. Joshi, Univ. of California, Los Angeles (United States)
- 7359 04 **Attosecond slicing of an LWFA produced electron beam (Invited Paper)** [7359-03]
C. M. S. Sears, A. Buck, K. Schmid, L. Veisz, D. Herrmann, J. Mikhailova, R. Tautz, F. Krausz, Max-Planck-Institut für Quantenoptik (Germany)
- 7359 05 **Coherent Thomson scattering at laser compressed and accelerated electron bunches (Invited Paper)** [7359-04]
D. an der Brügge, A. Pukhov, Heinrich-Heine-Univ. (Germany)

SESSION 2 TOWARDS A COMPACT X-FEL II

- 7359 07 **Pepper-pot emittance measurement of laser-plasma wakefield accelerated electrons (Invited Paper)** [7359-06]
R. P. Shanks, M. P. Anania, E. Brunetti, S. Cipiccia, B. Ersfeld, J. G. Gallacher, R. C. Issac, M. R. Islam, G. Vieux, G. H. Welsh, S. M. Wiggins, D. A. Jaroszynski, Univ. of Strathclyde (United Kingdom)
- 7359 08 **Linear and non-linear Thomson-scattering x-ray sources driven by conventionally and laser plasma accelerated electrons (Invited Paper)** [7359-07]
A. Debus, S. Bock, M. Bussmann, T. E. Cowan, A. Jochmann, T. Kluge, S. D. Kraft, R. Sauerbrey, K. Zeil, U. Schramm, Forschungszentrum Dresden-Rossendorf (Germany)

SESSION 3 FREQUENCY SHIFTING AND PHOTON ACCELERATION I

- 7359 09 **The flying mirror: future brightest x-ray and γ -ray source (Invited Paper)** [7359-08]
T. Zh. Esirkepov, S. V. Bulanov, M. Kando, A. S. Pirozhkov, Japan Atomic Energy Agency (Japan); A. G. Zhidkov, Central Research Institute of Electric Power Industry (Japan)
- 7359 0A **Photon Landau damping of relativistic plasma waves (Invited Paper)** [7359-09]
J. T. Mendonça, GoLP/Instituto Superior Técnico (Portugal)

SESSION 4 HIGH-HARMONIC GENERATION I

- 7359 0D **Factors influencing the temporal characteristics of coherent wake field harmonic emission from solid surfaces (Invited Paper)** [7359-12]
R. Hörlein, Max-Planck-Institut für Quantenoptik (Germany) and Ludwig-Maximilians-Univ. München (Germany); Y. Nomura, Max-Planck-Institut für Quantenoptik (Germany) and Univ. of Tokyo (Japan); S. G. Rykovanov, Max-Planck-Institut für Quantenoptik (Germany) and Moscow Physics Engineering Institute (Russian Federation); F. Krausz, Max-Planck-Institut für Quantenoptik (Germany) and Ludwig-Maximilians-Univ. München (Germany); G. D. Tsakiris, Max-Planck-Institut für Quantenoptik (Germany)
- 7359 0E **Attosecond and femtosecond metrology for plasma mirrors (Invited Paper)** [7359-13]
F. Quéré, H. George, Ph. Martin, CEA, IRAMIS, Service des Photons Atomes et Molécules (France)

SESSION 5 HIGH-HARMONIC GENERATION II

- 7359 0G **Ultrabright attosecond sources from relativistically oscillating mirrors (Invited Paper)** [7359-15]
M. Zepf, D. Adams, B. Dromey, M. Geissler, Queen's Univ. (United Kingdom); R. Hörlein, Y. Nomura, S. Rykovanov, G. D. Tsakiris, Max-Planck-Institut for Quantum Optics (Germany)

SESSION 6 FREQUENCY SHIFTING AND PHOTON ACCELERATION II

- 7359 0J **Short-wavelength magnetic structures from the plasma magnetic mode and their applications (Invited Paper)** [7359-18]
F. Fiuza, S. F. Martins, R. A. Fonseca, L. O. Silva, GoLP/Instituto de Plasma e Fusão Nuclear, Instituto Superior Técnico (Portugal)

7359 OK **XUV and IR electromagnetic radiation from nonlinear laser-plasma interaction (Invited Paper)** [7359-19]
M. Kando, A. S. Pirozhkov, Y. Fukuda, T. Zh. Esirkepov, I. Daito, K. Kawase, Japan Atomic Energy Agency (Japan); J. Ma, L. Chen, Japan Atomic Energy Agency (Japan) and Institute of Physics (China); Y. Hayashi, M. Mori, K. Ogura, H. Kotaki, A. Sagisaka, Japan Atomic Energy Agency (Japan); E. N. Ragozin, P. N. Lebedev Physical Institute (Russian Federation); A. Faenov, T. Pikuz, Japan Atomic Energy Agency (Japan) and Joint Institute for High Temperature (Russian Federation); H. Kiriya, H. Okada, T. Kameshima, J. K. Koga, T. Kawachi, H. Daido, Japan Atomic Energy Agency (Japan); T. Kimura, Japan Atomic Energy Agency (Japan) and Japan Synchrotron Radiation Research Institute (Japan); Y. Kato, Japan Atomic Energy Agency (Japan) and The Graduate School for the Creation of New Photonics Industries (Japan); T. Tajima, Japan Atomic Energy Agency (Japan) and Ludwig-Maximilians-Univ. München (Germany); S. V. Bulanov, Japan Atomic Energy Agency (Japan) and A. M. Prokhorov Institute of General Physics (Russian Federation)

7359 OL **Photon frequency up-shifting by an amplified plasma density wake due to two co-propagating laser pulses (Invited Paper)** [7359-20]
G. Raj, M. R. Islam, B. Ersfeld, D. A. Jaroszynski, Univ. of Strathclyde (United Kingdom)

SESSION 7 RAMAN AND BRILLOUIN AMPLIFICATION

7359 OM **Raman amplification in plasma: thermal effects and damping (Invited Paper)** [7359-21]
J. P. Farmer, B. Ersfeld, G. Raj, D. A. Jaroszynski, Univ. of Strathclyde (United Kingdom)

7359 ON **Experimental investigation of identical wavelength short light pulses crossing in underdense plasma (Invited Paper)** [7359-22]
L. Lancia, LULI, Ecole Polytechnique, CNRS, CEA, UPMC (France) and Univ. di Roma La Sapienza (France); J.-R. Marquès, J. Fuchs, M. Nakatsutsumi, A. Mancic, LULI, Ecole Polytechnique, CNRS, CEA, UPMC (France); P. Antici, Univ. di Roma La Sapienza (France); C. Riconda, LULI, Ecole Polytechnique, CNRS, CEA, UPMC (France); S. Weber, V. T. Tikhonchuk, CELIA, Univ. Bordeaux 1, CNRS, CEA (France); A. Héron, S. Hüller, J.-C. Adam, Ctr. de Physique Théorique, Ecole Polytechnique, CNRS (France); P. Audebert, LULI, Ecole Polytechnique, CNRS, CEA, UPMC (France)

7359 OO **Temporal evolution of density and field amplitudes in Raman amplification including relativistic and ponderomotive effects (Invited Paper)** [7359-23]
G. Raj, J. Farmer, B. Ersfeld, D. A. Jaroszynski, Univ. of Strathclyde (United Kingdom)

7359 OP **Effects of energy absorption on Raman amplification in plasma (Invited Paper)** [7359-24]
B. Ersfeld, J. Farmer, G. Raj, D. A. Jaroszynski, Univ. of Strathclyde (United Kingdom)

7359 OQ **Study of chirped pulse amplification based on Raman backscattering (Invited Paper)** [7359-25]
X. Yang, G. Vieux, A. Lyachev, J. Farmer, G. Raj, B. Ersfeld, E. Brunetti, M. Wiggins, R. Issac, D. A. Jaroszynski, Univ. of Strathclyde (United Kingdom)

7359 OR **Chirped pulse Raman amplification in plasma: high gain measurements (Invited Paper)** [7359-26]
G. Vieux, X. Yang, A. Lyachev, B. Ersfeld, J. Farmer, E. Brunetti, M. Wiggins, R. Issac, G. Raj, D. A. Jaroszynski, Univ. of Strathclyde (United Kingdom)

SESSION 8 BETATRON RADIATION I

- 7359 0T **Comparative study of betatron radiation from laser-wakefield and direct-laser accelerated bunches of relativistic electrons (Invited Paper)** [7359-28]
S. Kneip, Imperial College London (United Kingdom); C. McGuffey, Univ. of Michigan, Ann Arbor (United States); S. R. Nagel, C. Palmer, C. Bellei, J. Schreiber, Imperial College London (United Kingdom); C. Huntington, F. Dollar, T. Matsuoka, V. Chvykov, G. Kalintchenko, V. Yanovsky, A. Maksimchuk, University of Michigan, Ann Arbor (United States); K. Ta Phuoc, Lab. d'Optique Appliqué, ENSTA, Ecole Polytechnique (France); S. P. D. Mangles, Imperial College London (United Kingdom); K. Krushelnick, Imperial College London (United Kingdom) and University of Michigan, Ann Arbor (United States); Z. Najmudin, Imperial College London (United Kingdom)
- 7359 0V **Radiation post-processing in PIC codes (Invited Paper)** [7359-30]
J. L. Martins, S. F. Martins, R. A. Fonseca, L. O. Silva, GoLP/Instituto de Plasmas e Fusão Nuclear, Instituto Superior Técnico (Portugal)

SESSION 9 BETATRON RADIATION II

- 7359 0X **Temporal characterization and flux improvement of the x-ray betatron source (Invited Paper)** [7359-32]
R. Fitour, K. Ta Phuoc, S. Corde, A. Rousse, Lab. d'Optique Applique (France)

SESSION 10 TERAHERTZ SOURCES

- 7359 10 **Novel THz radiation from relativistic laser-plasmas (Invited Paper)** [7359-35]
Z. M. Sheng, Shanghai Jiao Tong Univ. (China) and Institute of Physics (China); H. C. Wu, W. M. Wang, X. G. Dong, M. Chen, Institute of Physics (China); J. Zhang, Shanghai Jiao Tong Univ. (China) and Institute of Physics (China)
- 7359 11 **Possibility of high power THz radiation via electromagnetically induced transparency at ion acoustic frequency region in laser-produced dense plasmas (Invited Paper)** [7359-36]
N. Yugami, T. Higashiguchi, Utsunomiya Univ. (Japan) and Japan Science and Technology Agency, CREST (Japan); R. Kodama, Osaka Univ. (Japan) and Japan Science and Technology Agency, CREST (Japan)
- 7359 12 **Terahertz radiation from the interaction of laser pulses with gas target (Invited Paper)** [7359-37]
M. Chen, A. Pukhov, X.-Y. Peng, O. Willi, Heinrich-Heine-Univ. Düsseldorf (Germany); Z.-M. Sheng, J. Zhang, Institute of Physics (China)

SESSION 11 BEAMS, PLASMA CHANNELS, AND DIAGNOSTICS

- 7359 14 **Narrow spread electron beams from a laser-plasma wakefield accelerator (Invited Paper)** [7359-39]
S. M. Wiggins, M. P. Anania, E. Brunetti, S. Cipiccia, B. Ersfeld, M. R. Islam, R. C. Issac, G. Raj, R. P. Shanks, G. Vieux, G. H. Welsh, Univ. of Strathclyde (United Kingdom); W. A. Gillespie, Univ. of Dundee (United Kingdom); A. M. MacLeod, Univ. of Abertay Dundee (United Kingdom); D. A. Jaroszynski, Univ. of Strathclyde (United Kingdom)

- 7359 15 **Electron beam pointing stability of a laser wakefield accelerator (Invited Paper)** [7359-40]
R. C. Issac, G. Vieux, G. H. Welsh, R. Shanks, E. Brunetti, S. Cipiccia, M. P. Anania, X. Yang,
S. M. Wiggins, M. R. Islam, B. Ersfeld, J. Farmer, G. Raj, S. Chen, D. Clark, T. McCanny,
D. A. Jaroszynski, University of Strathclyde (United Kingdom)

SESSION 12 TOWARDS A COMPACT X-FEL III

- 7359 16 **Transport of ultra-short electron bunches in a free-electron laser driven by a laser-plasma wakefield accelerator (Invited Paper)** [7359-41]
M. P. Anania, D. Clark, Univ. of Strathclyde (United Kingdom); S. B. van der Geer, M. J. de Loos, Technische Univ. Eindhoven (Netherlands); R. Isaac, A. J. W. Reitsma, G. H. Welsh, S. M. Wiggins, D. A. Jaroszynski, Univ. of Strathclyde (United Kingdom)

POSTER SESSION

- 7359 17 **Plasma diagnostics of a capillary plasma using pulse power (Invited Paper)** [7359-42]
T. Higashiguchi, Utsunomiya Univ. (Japan) and Japan Science and Technology Agency, CREST (Japan); H. Terauchi, Utsunomiya University (Japan); J. Bai, Utsunomiya Univ. (Japan); T. Kikuchi, Nagaoka Univ. of Technology (Japan); N. Yugami, Utsunomiya Univ. (Japan) and Japan Science and Technology Agency, CREST (Japan); T. Yatagai, Utsunomiya Univ. (Japan)
- 7359 18 **Compression of an ultrashort laser pulse via self-modulation in argon gas (Invited Paper)** [7359-43]
T. Higashiguchi, Utsunomiya Univ. (Japan) and Japan Science and Technology Agency, CREST (Japan); M. Kudo, Utsunomiya Univ. (Japan); N. Yugami, Utsunomiya Univ. (Japan) and Japan Science and Technology Agency, CREST (Japan); E. Takahashi, The Institute of Physical and Chemical Research (Japan)
- 7359 19 **Femto-second ultrashort laser wakefield electron bunch-duration measurements: a prism-based dispersion visible-to-IR spectrometer** [7359-45]
J. Lim, J. Faure, Lab. d'Optique Appliquée, ENSTA/CNRS, École Polytechnique (France); G. Gallot, Lab. d'Optique et Biosciences, CNRS, École Polytechnique-ENSTA (France); O. Lundh, C. Rechatin, V. Malka, Lab. d'Optique Appliquée, ENSTA/CNRS, École Polytechnique (France)

Author Index

Conference Committee

Symposium Chairs

Pavel Tomanek, Brno University of Technology (Czech Republic)
Alan G. Michette, King's College London (United Kingdom)
Bahaa E. A. Saleh, Boston University (United States)

Symposium Honorary Chair

Jan Perina, Sr., Palacký University (Czech Republic)

Conference Chairs

Dino A. Jaroszynski, University of Strathclyde (United Kingdom)
Antoine Rouse, Ecole Nationale Supérieure de Techniques Avancées
(France)

Program Committee

Robert Bingham, Science and Technology Facilities Council (United Kingdom)
Christopher E. Clayton, University of California, Los Angeles (United States)
N. Piovela, Università degli studi di Milano (Italy)
Alexander Pukhov, Heinrich-Heine-Universität Düsseldorf (Germany)
Luca Serafini, Università degli studi di Milano (Italy)
Gennady Shvets, The University of Texas at Austin (United States)
Luis O. Silva, Instituto Superior Técnico (Portugal)
Mark S. Wiggins, University of Strathclyde (United Kingdom)
Matthew Zepf, Queen's University Belfast (United Kingdom)

Session Chairs

- 1 Towards a Compact X-FEL I
Luis O. Silva, Instituto Superior Técnico (Portugal)
- 2 Towards a Compact X-FEL II
Christopher E. Clayton, University of California, Los Angeles (United States)
- 3 Frequency Shifting and Photon Acceleration I
Christopher E. Clayton, University of California, Los Angeles (United States)

- 4 High-Harmonic Generation I
Mark S. Wiggins, University of Strathclyde (United Kingdom)
- 5 High-Harmonic Generation II
Tito Mendonca, Instituto Superior Técnico (Portugal)
- 6 Frequency Shifting and Photon Acceleration II
Tito Mendonca, Instituto Superior Técnico (Portugal)
- 7 Raman and Brillouin Amplification
Dino A. Jaroszynski, University of Strathclyde (United Kingdom)
- 8 Betatron Radiation I
Antoine Rousse, Ecole Nationale Supérieure de Techniques Avancées (France)
- 9 Betatron Radiation II
Alexander Pukhov, Heinrich-Heine-Universität Düsseldorf (Germany)
- 10 Terahertz Sources
Matthew Zepf, Queen's University Belfast (United Kingdom)
- 11 Beams, Plasma Channels, and Diagnostics
Matthew Zepf, Queen's University Belfast (United Kingdom)
- 12 Towards a Compact X-FEL III
Matthew Zepf, Queen's University Belfast (United Kingdom)

PHOTON PHYSICS: FROM WAVE MECHANICS TO QUANTUM ELECTRODYNAMICS

Ole Keller

Institute of Physics and Nanotechnology, Aalborg University, Denmark

ABSTRACT

When rewritten in an appropriate manner, the microscopic Maxwell-Lorentz equations appear as a wave-mechanical theory for photons, and their quantum physical interaction with matter. A natural extension leads from photon wave mechanics to quantum electrodynamics (QED). In its modern formulation photon wave mechanics has given us valuable new insight in subjects such as spatial photon localization, near-field photon dynamics, transverse photon mass, photon eikonal theory, photon tunneling, and rim-zone electrodynamics. The present review is based on my plenary lecture at the SPIE-Europe 2009 Optics and Optoelectronics International Symposium in Prague.

1. INTRODUCTION

The birth of quantum field radiation theory in the years 1925-30, did not cause physicists to give up the idea of a particle concept of light. Beautiful wave mechanical theories for the photon were thus established by Landau and Peierls in 1930,¹ and by Oppenheimer² the year after. For a brief historical review of photon wave mechanics the reader may consult reference three.³ Over the years physicists interest in photon wave mechanics, and its extension to the field-quantized level, has waxed and waned, but never fallen to rest.⁴ In the present review, I look at this almost century old field in a modern perspective, and point out some of the modern perspectives for low-energy physical optics.

It has been known for almost two centuries that the eikonal equation in geometrical optics is form-identical to the Hamilton-Jacobi equation for Hamilton's characteristic function in Newtonian mechanics.^{5,6} This fact led almost exactly a century later De Broglie and Schrödinger to the wave mechanics of massive particles.^{3,5,6} In this perspective, it is perhaps not surprising that classical electrodynamics, based on the microscopic Maxwell-Lorentz equations, may be looked upon as the wave-mechanical theory of the photon. Starting from the Klein-Gordon equation, the quantum-mechanical eikonal equation for a relativistic spinless particle subjected to an electromagnetic field is first established.

In free space various constructions (objects) can claim to represent a photon wave function. Here, we discuss two of the most attractive suggestions: the photon energy wave function,^{2,4} and a wave function based on the transverse vector potential.⁷ For the potential approach, we set up the photon Schrödinger equation in both momentum and configuration space. Afterwards, we discuss the transverse photon mass concept in diamagnetic field-matter interaction, and the photon eikonal theory.

Modern photon wave mechanics appears to be particularly important in the quantum theory of near-field electrodynamics.^{8,9} In the rim zone of matter virtual photons play a significant role, but only in light-matter interactions. A (new) virtual photon type called a near-field photon is introduced and its wave mechanics analyzed. The near-field photon is accompanied by a gauge photon, and the two photons place in QED is discussed. I end the review by a brief discussion of the position operator problem for massive photons (plasmaitons), and emphasize the importance of the rim zone in treatments of the spatial localization problem for photons emitted from microscopic (or mesoscopic) sources.^{10,11}

2. GEOMETRICAL OPTICS AND NEWTONIAN MECHANICS IN THE HAMILTON-JACOBI FORMULATION

It has been known for almost two centuries that the dynamics of a system of interacting classical (Newtonian) particles can be described neatly on the basis of Hamilton's equations

$$\dot{q}_i = \frac{\partial}{\partial p_i} H(\{q\}, \{p\}, t) \quad (1)$$

$$\dot{p}_i = -\frac{\partial}{\partial q_i} H(\{q\}, \{p\}, t) \quad (2)$$

where q_i is the generalized coordinate number i , and p_i its associated canonical (conjugate) momentum. The dots stand for differentiation with respect to time. As indicated, the Hamilton function H depends in general on the full set of coordinates and momenta, and on time. By a particular canonical transformation, which generating function involves Hamilton's principal function $S(\{q\}, \{P\}, t)$, the new Hamiltonian can be required to vanish identically. This ensures that the new generalized coordinates and momenta ($\{P\}$) become *time-independent*, and can be taken as the initial values of the old phase space variables. Hamilton's principal function now must satisfy the partial differential equation (called the Hamilton-Jacobi equation)

$$H\left(\{q\}, \left\{\frac{\partial S}{\partial q}\right\}, t\right) + \frac{\partial S}{\partial t} = 0, \quad (3)$$

and the (old) generalized momenta are identified as

$$p_i = \frac{\partial}{\partial q_i} S(\{q\}, t), \quad (4)$$

leaving out the reference to $\{P\}$ from the notation. In the special but important case where H does not depend explicitly on time, a trial solution of the form

$$S(\{q\}, t) = W(\{q\}) - Et \quad (5)$$

leads to the Hamilton-Jacobi equation

$$H\left(\{q\}, \left\{\frac{\partial W}{\partial q}\right\}\right) = E \quad (6)$$

for Hamilton's characteristic function, $W(\{q\})$. Usually, the constant E represents the (conserved) energy of the system. For a single particle in a time-independent potential, $V(\mathbf{r})$, the equation for Hamilton's characteristic function takes the following form:

$$\nabla W(\mathbf{r}) \cdot \nabla W(\mathbf{r}) = 2m(E - V(\mathbf{r})), \quad (7)$$

where m is the particle's mass, and \mathbf{r} its position vector. With the identification in Eq. (4), which here on vector form gives $\mathbf{p} = \nabla W$, Eq. (7) expresses the fact that the particle's total energy is the sum of the kinetic and potential energies.

Geometrical optics deals with the phenomena for which the wavelength of light can be neglected. In this short-wavelength limit ($\lambda \rightarrow 0$) the optical laws may be formulated in the language of geometry, and the basic quantity is the so-called eikonal $S(\mathbf{r})$, which enters the various (monochromatic) electromagnetic field vectors (\mathbf{V}) via the ansatz $\mathbf{V}(\mathbf{r}) = \mathbf{V}_0 \exp(iq_0 S(\mathbf{r}))$, $q_0 = \omega/c_0$ being the vacuum wavenumber of light at the angular frequency ω . The fundamental equations of geometrical optics is the eikonal equation⁶

$$\nabla S(\mathbf{r}) \cdot \nabla S(\mathbf{r}) = n^2(\mathbf{r}), \quad (8)$$

where $n(\mathbf{r})$ is the refractive index, which in the lowest-order approximation is taken as a real quantity. As first noticed by Hamilton in 1827 there is a striking form-identity between the characteristic function [Eq. (7)] and the eikonal equation of geometrical optics [Eq. (8)]. Classical electrodynamics based in the macroscopic Maxwell equations (or in a somewhat broader perspective the microscopic Maxwell-Lorentz equations) is the covering

theory of geometrical optics, and this fact led almost exactly one hundred years later De Broglie and Schrödinger to the covering theory of Newtonian mechanics, i.e., quantum mechanics, as might be well-known to the reader.

For what follows it is valuable briefly to consider the quantum mechanical eikonal theory for a relativistic spinless particle of rest mass m and charge q . Placed in an electromagnetic field described by the contravariant four-potential $\{A^\mu\} = (A^0 = \Phi/c_0, \mathbf{A})$ the classical relativistic Hamiltonian is

$$H = c_0 [(mc_0)^2 + (\mathbf{p} - q\mathbf{A})^2]^{1/2} + q\Phi. \quad (9)$$

The Hamiltonian in Eq. (9) has no explicit time dependence, and with $\mathbf{p} = \nabla S$, the equation for Hamilton's principal function [Eq. (3)] can be written down. All solutions to this equation will also be solutions to the squared Hamilton-Jacobi equation, viz.,

$$c_0^2 (\nabla S - q\mathbf{A})^2 + (mc_0^2)^2 = \left(\frac{\partial S}{\partial t} + q\Phi \right)^2. \quad (10)$$

The quantum mechanical eikonal equation for a relativistic spinless particle subjected to an electromagnetic field is obtained starting from the Klein-Gordon equation

$$c_0^2 \left[\left(\frac{\hbar}{i} \nabla - q\mathbf{A} \right)^2 + (mc_0)^2 \right] \Psi(\mathbf{r}, t) = \left(\frac{\hbar}{i} \frac{\partial}{\partial t} + q\Phi \right)^2 \Psi(\mathbf{r}, t). \quad (11)$$

By inserting the eikonal ansatz

$$\Psi(\mathbf{r}, t) = \Psi_0 \exp \left(\frac{i}{\hbar} S(\mathbf{r}, t) \right) \quad (12)$$

for the wavefunction, the quantum mechanical Hamilton-Jacobi equation takes the form

$$(\nabla S - q\mathbf{A})^2 + (mc_0)^2 - \frac{1}{c_0^2} \left(\frac{\partial S}{\partial t} + q\Phi \right)^2 = i\hbar \left(\nabla^2 - \frac{1}{c_0^2} \frac{\partial^2}{\partial t^2} \right) S \quad (13)$$

in the Lorenz gauge. In the classical limit ($\hbar \rightarrow 0$), Eq. (13) is reduced to Eq. (10). In covariant notation, one has in general

$$(\partial_\mu S - qA_\mu)(\partial^\mu S - qA^\mu) + (mc_0)^2 = i\hbar \partial_\mu (\partial^\mu S - qA^\mu), \quad (14)$$

and with the Lorenz gauge condition, given by

$$\partial_\mu A^\mu = 0, \quad (15)$$

Eq. (14) is reduced to the form given in Eq. (13). The relativistic covariance of the eikonal equation is manifest when the equation is written in the form given in Eq. (14).

3. PHOTON WAVE MECHANICS

The analysis in section 2 suggests that the microscopic Maxwell-Lorentz equations somehow may be considered as a wave-mechanical theory of the photon. This possibility was originally investigated by Landau and Peierls¹ and Oppenheimer,² and over the years physicist's interest in photon wave mechanics has waxed and waned, but never fallen to rest. In free space, various "constructions" (objects) can rightly claim that they represent the wave function of the photon. What is important physically is that the observable predictions related to the photon-matter interaction will be identical for the various choices.

In the energy wave-function formalism two photon wave functions, $\mathbf{f}_\pm^{(+)}$, are defined as follows:⁴

$$\mathbf{f}_\pm^{(+)}(\mathbf{r}, t) = \left(\frac{\epsilon_0}{2} \right)^{1/2} \left[\mathbf{e}_T^{(+)}(\mathbf{r}, t) \pm ic_0 \mathbf{b}^{(+)}(\mathbf{r}, t) \right]. \quad (16)$$

The superscript (+) indicates that the wave function is build from positive-frequency components of the electromagnetic field. In the language of coherence theory the $\mathbf{f}_\pm^{(+)}$'s are analytical signals. The wave function of

the antiphoton is constructed from the negative-frequency part of the field spectrum. Physically, the photon and antiphoton are identical. This can be inferred from the Maxwell equations. Small letters \mathbf{e}_T and \mathbf{b} have been assigned to the (analytical) electric and magnetic fields of the photon to indicate that the wave function has been properly normalized. To normalize the wave function of the in general non-monochromatic photon it is necessary to adjust the amplitude of the current-density source distribution correctly.⁸ To emphasize that the electrical field in free space is a transverse vector field, a subscript T has been added to $\mathbf{e}^{(+)}$. As we shall understand very soon, $\mathbf{f}_+^{(+)}$ and $\mathbf{f}_-^{(+)}$ are wave functions composed of positive- and negative-helicity components. The photon helicity is a Lorentz-invariant property.

From the free-space Maxwell equations one can establish the following quantum mechanical wave equations for the two helicity species:

$$i\hbar \frac{\partial}{\partial t} \mathbf{f}_{\pm}^{(+)}(\mathbf{r}, t) = \pm c_0 \left(\frac{\mathbf{s}}{\hbar} \cdot \frac{\hbar}{i} \nabla \right) \mathbf{f}_{\pm}^{(+)}(\mathbf{r}, t). \quad (17)$$

In Eq. (17) we have written the wave equation in Hamiltonian form. The vector \mathbf{s} is the Cartesian spin-one operator of the photon. In a 3×3 matrix representation the Cartesian components of \mathbf{s} are given by

$$(s_k)_{ij} = \frac{\hbar}{i} \epsilon_{ijk}, \quad (18)$$

where ϵ_{ijk} is the Levi-Civita tensor. Since $\hat{\mathbf{p}} = (\hbar/i)\nabla$ is the photon momentum operator in configuration space, the scalar product $\mathbf{s} \cdot \hat{\mathbf{p}}$ confirms that the wave functions $\mathbf{f}_{\pm}^{(+)}(\mathbf{r}, t)$ describe photons composed of positive- and negative helicity components. Since Planck's constant (\hbar) does not appear in the classical Maxwell-Lorentz equations, \hbar can of course be eliminated from Eq. (17), as the reader readily may realize. The name photon energy wave function relates to the fact that the quantum mechanical mean value of the photon energy-momentum four vector operator $\{p^\mu\} = (\hbar\omega/c_0, \hbar\mathbf{q})$ is given by

$$\{P^\mu\} = \frac{1}{\hbar c_0} \sum_{s=+,-} \int_{-\infty}^{\infty} \{p^\mu\} |f_s^{(+)}(\mathbf{q}; t)|^2 \frac{d^3q}{(2\pi)^3 q}, \quad (19)$$

where the $f_{\pm}^{(+)}(\mathbf{q}; t)$'s are the scalar photon wave functions in momentum space.

The vectorial and scalar wave functions are related by $\mathbf{f}_{\pm}^{(+)}(\mathbf{q}; t) = f_{\pm}^{(+)}(\mathbf{q}; t) \mathbf{e}_{\pm}(\mathbf{q}/q)$, where \mathbf{e}_{\pm} are the helicity unit vectors of the + and - species. Relativistically, d^3q/q is the invariant volume element on the light cone.

It is often convenient to use spinor notation, and thus combine the two wave functions in Eq. (16) into a single one, viz.,

$$\Phi(\mathbf{r}, t) = \begin{pmatrix} \mathbf{f}_+^{(+)}(\mathbf{r}, t) \\ \mathbf{f}_-^{(+)}(\mathbf{r}, t) \end{pmatrix}. \quad (20)$$

The photon wave function in Eq. (20) is a six-component object.

In the present context it is useful to base photon wave mechanics on the electromagnetic potentials. In free space only a transverse vector potential exists, and this potential is gauge invariant. The analytical part of the transverse vector potential $\mathbf{a}_T^{(+)}(\mathbf{r}, t)$ satisfies the wave equation

$$\square \mathbf{a}_T^{(+)}(\mathbf{r}, t) = \mathbf{0}, \quad (21)$$

where $\square = \nabla^2 - c_0^{-2} \partial^2 / \partial t^2$ is the d'Alembertian operator. Formally, one may factorize the \square -operator as follows:

$$\square = \left(\frac{i}{c_0} \frac{\partial}{\partial t} + \sqrt{-\nabla^2} \right) \left(\frac{i}{c_0} \frac{\partial}{\partial t} - \sqrt{-\nabla^2} \right), \quad (22)$$

and from this it appears that all solutions to

$$i\hbar \frac{\partial}{\partial t} \mathbf{a}_T^{(+)}(\mathbf{r}, t) = \hbar c_0 \sqrt{-\nabla^2} \mathbf{a}_T^{(+)}(\mathbf{r}, t) \quad (23)$$

are also solutions to Eq. (21). One may consider $\mathbf{a}_T^{(+)}(\mathbf{r}, t)$ [small letter for normalization] as the vectorial wave function for the photon in free space, and Eq. (23) as its quantum mechanical wave equation, written in Hamiltonian form. The quantity $\sqrt{-\nabla^2}$ is a spatially nonlocal operator which action in Fourier (wave-vector) space is given via

$$\sqrt{-\nabla^2}\mathbf{a}_T^{(+)}(\mathbf{r}, t) = \int_{-\infty}^{\infty} q\mathbf{a}_T^{(+)}(\mathbf{q}; t)e^{i\mathbf{q}\cdot\mathbf{r}}\frac{d^3q}{(2\pi)^3}. \quad (24)$$

The operator $i\sqrt{-\nabla^2}$ was originally introduced by Landau and Peierls in 1930 in connection with their attempt to establish a wave equation for the light quantum (photon). In \mathbf{q} -space, Eq. (23) takes the form

$$i\hbar\frac{\partial}{\partial t}\mathbf{a}_T^{(+)}(\mathbf{q}; t) = c_0\hbar q\mathbf{a}_T^{(+)}(\mathbf{q}; t), \quad (25)$$

and this equation may be considered as the vectorial wave equation for a transverse photon with momentum $\mathbf{p} = \hbar\mathbf{q}$ and energy $H = pc_0$. Two transverse scalar photon wave functions can be introduced in the $(\mathbf{q}; t)$ -domain by expanding $\mathbf{a}_T^{(+)}(\mathbf{q}; t)$ after two orthogonal (possibly complex) unit vectors (helicity unit vectors, e.g) located in a plane perpendicular to \mathbf{q} .

4. TRANSVERSE PHOTON MASS. PHOTON EIKONAL THEORY

Let us now consider the photon propagation in a solid-state plasma (jellium), and let us limit ourselves to high frequencies and linear photon-matter interactions. At high frequencies the particle properties of the photon dominates over its wave properties, and the field-matter coupling is diamagnetic. In the space-frequency domain the analytical part of the transverse vectorial potential hence satisfies the integro-differential equation¹²

$$(\nabla^2 + q_0^2)\mathbf{a}_T^{(+)}(\mathbf{r}; \omega) = \frac{\mu_0 e^2}{m} \int_{-\infty}^{\infty} \overset{\leftrightarrow}{\delta}_T(\mathbf{r} - \mathbf{r}') \cdot \left[N(\mathbf{r}')\mathbf{a}_T^{(+)}(\mathbf{r}'; \omega) \right] d^3r', \quad (26)$$

where $N(\mathbf{r})$ is the many-body electron density, $\overset{\leftrightarrow}{\delta}_T(\mathbf{r} - \mathbf{r}')$ is the transverse delta function, $q_0 = \omega/c_0$ is the vacuum wave number of light, and e and m the electron charge and mass. In the present context there is a particularly interesting special case of Eq. (26). Thus, if the electron density variations in space are negligible [$N(\mathbf{r}) \cong N_0$] it can be shown that $\mathbf{a}_T^{(+)}(\mathbf{r}, t)$ satisfies the equation

$$i\hbar\frac{\partial}{\partial t}\mathbf{a}_T^{(+)}(\mathbf{r}, t) = c_0\hbar\sqrt{Q_c^2 - \nabla^2}\mathbf{a}_T^{(+)}(\mathbf{r}, t), \quad (27)$$

where the operator $(Q_c^2 - \nabla^2)^{1/2}$ acts as follows in Fourier space:

$$\sqrt{Q_c^2 - \nabla^2}\mathbf{a}_T^{(+)}(\mathbf{r}, t) = \int_{-\infty}^{\infty} \sqrt{Q_c^2 + q^2}\mathbf{a}_T^{(+)}(\mathbf{q}; t)e^{i\mathbf{q}\cdot\mathbf{r}}\frac{d^3q}{(2\pi)^3}. \quad (28)$$

The quantity

$$Q_c = \frac{1}{c_0} \left(\frac{N_0 e^2}{m\epsilon_0} \right)^{1/2} \quad (29)$$

is the electron system's plasma wave number. One may consider Eq. (27) as the quantum mechanical wave equation for the transverse photon (plasmariton). In the wave-vector-time domain Eq. (27) takes the form

$$i\hbar\frac{\partial}{\partial t}\mathbf{a}_T^{(+)}(\mathbf{q}; t) = c_0\hbar(q^2 + Q_c^2)^{1/2}\mathbf{a}_T^{(+)}(\mathbf{q}; t), \quad (30)$$

and from this one may show that the energy-momentum relation for the plasmariton takes the relativistic form

$$E = + [(pc_0)^2 + (Mc_0^2)^2]^{1/2}. \quad (31)$$

In a sense one may now claim that the transverse photon has acquired a finite mass

$$M = \frac{\hbar Q_c}{c_0} \quad (32)$$

in its high-frequency interaction with the jellium. The quantity $Q_c = Mc_0/\hbar$ is the Compton wave number of the massive transverse photon.

The wave equation for the plasmariton is form-identical to the Klein-Gordon equation, and the eikonal ansatz $\mathbf{a}_T^{(+)}(\mathbf{r}, t) = \mathbf{a}_{T,0}^{(+)} \exp(iS(\mathbf{r}, t)/\hbar)$ therefore leads to the following eikonal equation for the "free" massive photon:

$$(\partial_\mu S)(\partial^\mu S) + (Mc_0)^2 = i\hbar\partial_\mu\partial^\mu S. \quad (33)$$

If the spatial variations in $N(\mathbf{r})$ is taken into account the eikonal problem becomes more complicated.

5. THE NEAR-FIELD PHOTON CONCEPT

In near-field electrodynamics one must include two other types of photons, viz., longitudinal and scalar photons. These so-called virtual photons, which do not exist as free particles, we hence characterize by the analytical vector potentials $a_L^{(+)}(\mathbf{q}; t)[\mathbf{a}_L^{(+)} = a_L^{(+)}\mathbf{q}/q]$ and $a_0^{(+)}(\mathbf{q}; t)$. Instead of using the longitudinal and scalar photon variables it is convenient in near-field optics to introduce two new photon types via the unitary transformation⁹

$$\begin{pmatrix} a_{NF}^{(+)}(\mathbf{q}; t) \\ a_G^{(+)}(\mathbf{q}; t) \end{pmatrix} = \frac{1}{\sqrt{2}} \begin{pmatrix} i & -i \\ 1 & 1 \end{pmatrix} \begin{pmatrix} a_L^{(+)}(\mathbf{q}; t) \\ a_0^{(+)}(\mathbf{q}; t) \end{pmatrix}. \quad (34)$$

The quantities $a_{NF}^{(+)}(\mathbf{q}; t)$ and $a_G^{(+)}(\mathbf{q}; t)$ are the quantum mechanical wave functions of the near-field (NF) and gauge (G) photons. The theory for $a_L^{(+)}$ and $a_0^{(+)}$, or equivalently $a_{NF}^{(+)}$ and $a_G^{(+)}$, always is formulated in the Lorenz gauge. The primary quantity in near-field electrodynamics is the analytical longitudinal electric field $\mathbf{E}_L^{(+)}(\mathbf{q}; t) = E_L^{(+)}(\mathbf{q}; t)\mathbf{q}/q$. It may be shown that the Lorenz gauge condition implies that $E_L^{(+)}(\mathbf{q}; t)$ can be expressed solely in terms of either $a_{NF}^{(+)}(\mathbf{q}; t)$ or $a_G^{(+)}(\mathbf{q}; t)$, i.e.,

$$E_L^{(+)}(\mathbf{q}; t) = \sqrt{2} \left(c_0q + i\frac{\partial}{\partial t} \right) a_{NF}^{(+)}(\mathbf{q}; t) = -i\sqrt{2} \left(c_0q - i\frac{\partial}{\partial t} \right) a_G^{(+)}(\mathbf{q}; t). \quad (35)$$

In free space the near-field and gauge photon wave functions satisfy the Hamiltonian wave equations

$$i\hbar\frac{\partial}{\partial t}a_{NF}^{(+)}(\mathbf{q}; t) = c_0\hbar qa_{NF}^{(+)}(\mathbf{q}; t), \quad (36)$$

$$i\hbar\frac{\partial}{\partial t}a_G^{(+)}(\mathbf{q}; t) = c_0\hbar qa_G^{(+)}(\mathbf{q}; t), \quad (37)$$

and since $E_L(\mathbf{q}; t) = 0$ in free space, Eqs. (35) and (36) imply that $a_{NF}^{(+)}(\mathbf{q}; t) = 0$. One cannot conclude from these equations that the wave function of the gauge photon vanishes in free space. A gauge transformation to a new (NEW) gauge within the Lorenz gauge can make $a_G^{(+)}{}^{NEW}(\mathbf{q}; t) = 0$. The name gauge photon originates in this property. In the presence of matter, the near-field and gauge photon variables satisfy the dynamical equations

$$\left(c_0q + i\frac{\partial}{\partial t} \right) a_{NF}^{(+)}(\mathbf{q}; t) = \frac{1}{i\sqrt{2}\epsilon_0} \frac{1}{qc_0} J_0^{(+)}(\mathbf{q}; t), \quad (38)$$

$$\left(c_0q - i\frac{\partial}{\partial t} \right) a_G^{(+)}(\mathbf{q}; t) = \frac{1}{\sqrt{2}\epsilon_0} \frac{1}{qc_0} J_0^{(+)}(\mathbf{q}; t), \quad (39)$$

where $J_0^{(+)}(\mathbf{q}; t) = c_0\rho(\mathbf{q}; t)$ is the $\mu = 0$ component of the contravariant current density four-vector.

6. RIM ZONE ELECTRODYNAMICS

The near-field photon concept connects to optical near-field interactions via Eq. (35) and (38). I shall not here undertake a systematic analysis of this connection. The longitudinal part of the electric field, \mathbf{E}_L , plays an important role in near-field optics, and this field relates to the near-field photon variable via the analytical relation

$$\mathbf{E}_L^{(+)}(\mathbf{r}, t) = c_0 \sqrt{2} \left(\frac{i}{c_0} \frac{\partial}{\partial t} + \sqrt{-\nabla^2} \right) \mathbf{a}_{NF}^{(+)}(\mathbf{r}, t), \quad (40)$$

where $\mathbf{a}_{NF}^{(+)}(\mathbf{r}, t)$ is the spatial Fourier transform of the vectorial quantity $a_{NF}^{(+)}(\mathbf{q}; t)\mathbf{q}/q$. As an immediate consequence of Eqs. (35) and (38) the reader may obtain the following relation between the longitudinal electric field and the longitudinal source current density (\mathbf{J}_L):

$$\mathbf{J}_L(\mathbf{r}; \omega) = i\epsilon_0 \omega \mathbf{E}_L(\mathbf{r}; \omega). \quad (41)$$

In the presence of a transverse current density source, $\mathbf{J}_T^{(+)}(\mathbf{r}, t)$, the transverse photon variable satisfies the inhomogeneous wave equation

$$\square \mathbf{a}_T^{(+)}(\mathbf{r}, t) = -\mu_0 \mathbf{J}_T^{(+)}(\mathbf{r}, t), \quad (42)$$

and from this one can obtain the integral relation

$$\mathbf{E}_T(\mathbf{r}; \omega) = i\mu_0 \omega \int_{-\infty}^{\infty} g(|\mathbf{r} - \mathbf{r}'|; \omega) \mathbf{J}_T(\mathbf{r}'; \omega) d^3 r' \quad (43)$$

between the transverse parts of the electric field and source current density. The quantity $g(R; \omega) = (4\pi R)^{-1} \times \exp(iq_0 R)$ is the outgoing scalar propagator. The relations in Eqs. (41) and (43) form the standard starting point for classical near-field electrodynamics. Everywhere in space outside matter one has $\mathbf{J}_L = -\mathbf{J}_T$, since $\mathbf{J} = \mathbf{0}$. It appears from the equation

$$\mathbf{J}_L(\mathbf{r}; \omega) = \frac{1}{3} \mathbf{J}(\mathbf{r}; \omega) + \frac{1}{4\pi} PV \int_{-\infty}^{\infty} R^{-3} \left(\overleftrightarrow{\mathbf{U}} - 3 \frac{\mathbf{R}\mathbf{R}}{R^2} \right) \cdot \mathbf{J}(\mathbf{r}'; \omega) d^3 r', \quad (44)$$

where $\mathbf{R} = \mathbf{r} - \mathbf{r}'$, and PV stands for principal value that \mathbf{J}_L is appreciably different from zero only in a narrow zone outside a matter-filled region. This zone I have called the rim zone, and in this zone the longitudinal electric field is nonvanishing, and satisfies

$$\nabla \cdot \mathbf{E}_L(\mathbf{r}; \omega) = \nabla \times \mathbf{E}_L(\mathbf{r}; \omega) = 0, \quad (45)$$

the last equation per definition, of course. In the language of photon wave mechanics near-field optics describes how transverse photons via their interaction with near-field photons are generated and destroyed in space-time.

7. NEAR-FIELD AND GAUGE PHOTONS IN QED

The wave mechanical formalism can be extended to the quantum-electrodynamic level by the standard covariant quantization with an indefinite metric (Gupta-Bleuler formalism),^{13, 14} The Hamilton operator of the field becomes (in box quantization)

$$\hat{H} = \sum_{\mathbf{q}, r} \hbar \omega_q \hat{a}_r^\dagger(\mathbf{q}) \hat{a}_r(\mathbf{q}), \quad (46)$$

where $\omega_q = c_0 q$. The r-summation in Eq. (46) is over the two transverse modes (T1, T2), and the longitudinal (L) and scalar (0) modes. The annihilation ($\hat{a}_r(\mathbf{q})$) and creation ($\hat{a}_r^\dagger(\mathbf{q})$) operators of the various modes satisfy the commutation relations

$$[\hat{a}_r(\mathbf{q}), \hat{a}_s^\dagger(\mathbf{q}')] = \delta_{rs} \delta_{\mathbf{q}, \mathbf{q}'}. \quad (47)$$

By extension of Eq. (34) to the operator level it is easy to show that

$$\hat{a}_L^\dagger \hat{a}_L + \hat{a}_0^\dagger \hat{a}_0 = \hat{a}_{NF}^\dagger \hat{a}_{NF} + \hat{a}_G^\dagger \hat{a}_G, \quad (48)$$

a relation which allows one to eliminate the longitudinal and scalar photon operators in the field Hamiltonian in favour of the near-field and gauge photon operators. The annihilation and creation operators of these virtual photon types satisfy the commutator relations

$$\left[\hat{a}_{NF}(\mathbf{q}), \hat{a}_{NF}^\dagger(\mathbf{q}') \right] = \delta_{\mathbf{q}, \mathbf{q}'}, \quad (49)$$

$$\left[\hat{a}_G(\mathbf{q}), \hat{a}_G^\dagger(\mathbf{q}') \right] = \delta_{\mathbf{q}, \mathbf{q}'}, \quad (50)$$

and the NF-G commutators vanish. In free space the quantum Lorenz condition leads for all \mathbf{q} to

$$[\hat{a}_L(\mathbf{q}) - \hat{a}_0(\mathbf{q})] |\psi\rangle = \hat{a}_{NF}(\mathbf{q}) |\psi\rangle = 0, \quad (51)$$

so that no physical states $|\psi\rangle$ have near-field photons. The Lorenz condition puts no restrictions on the number of gauge photons in free space. Two different gauges within the Lorenz gauge correspond to two different excitations of G-modes, but as in the classical theory the gauge photon plays no physical role in free space.

8. PLASMARITON POSITION OPERATOR. SPATIAL PHOTON LOCALIZATION

Let us now ask the following question: to what extent can a point particle in the form of a transverse (massive) photon be localized in space? If a particle's position is to be measured in quantum mechanics, the various positions in space, \mathbf{r}_0 , should be the eigenvalues of some position operator (observable), $\hat{\mathbf{r}}$. To pin down elements of the problem, we first consider a massive relativistic particle of zero spin. For such a particle the Lorentz-invariant scalar product is given by¹⁵

$$\langle \psi | \phi \rangle = \int_{-\infty}^{\infty} \psi^*(\mathbf{p}) \phi(\mathbf{p}) \frac{d^3 p}{[p^2 + (Mc_0)^2]^{1/2}} \quad (52)$$

in the momentum representation. In the plasmariton case the particle mass (M) is given by Eq. (32). Within the framework of the relativistic scalar product the Hermitian operator

$$\hat{\mathbf{r}} = i\hbar \left\{ \nabla_{\mathbf{p}} - \frac{\mathbf{p}}{2[p^2 + (Mc_0)^2]} \right\} \quad (53)$$

turns out to be an acceptable position operator for a Klein-Gordon particle. In the non-relativistic (NR) limit one regains the naive position operator $\hat{\mathbf{r}}_{NR} = i\hbar \nabla_{\mathbf{p}}$. The operator in Eq. (53) satisfies the basic dyadic commutator relation

$$[\hat{\mathbf{r}}, \hat{\mathbf{p}}] = i\hbar \hat{\mathbf{1}} \quad (54)$$

where $\hat{\mathbf{1}}$ is the unit matrix operator. In the momentum representation, where the eigenvalue equation reads

$$\hat{\mathbf{r}}(\mathbf{p}) \psi_{\mathbf{r}_0}(\mathbf{p}) = \mathbf{r}_0 \psi_{\mathbf{r}_0}(\mathbf{p}), \quad (55)$$

the (unnormalizable) eigenfunction belonging to the eigenvalue \mathbf{r}_0 is given by

$$\psi_{\mathbf{r}_0}(\mathbf{p}) = [p^2 + (Mc_0)^2]^{1/4} \exp\left(-\frac{i}{\hbar} \mathbf{r}_0 \cdot \mathbf{p}\right), \quad (56)$$

apart from an arbitrary multiplicative factor. In *configuration space* the eigenfunction takes the form

$$\psi_{\mathbf{r}_0}(\mathbf{r}) = 2(2\pi\hbar)^{-2} \frac{(Mc_0)^{3/2}}{|\mathbf{r} - \mathbf{r}_0|} F(Q_c |\mathbf{r} - \mathbf{r}_0|), \quad (57)$$

where

$$F(Q_c |\mathbf{r} - \mathbf{r}_0|) = \int_0^\infty x(1+x^2)^{1/4} \sin(Q_c |\mathbf{r} - \mathbf{r}_0| x) dx. \quad (58)$$

It appears that the function F only depends on the ratio between $|\mathbf{r} - \mathbf{r}_0|$ and the Compton wavelength of the plasmariton, $\lambda_c = h/(Mc_0)$. Since $F(z) \approx z^{-5/4} \exp(-z)$ for $z \gg 1$, the wave function of a scalar particle

definitely localized at \mathbf{r}_0 is smeared out in configuration space over a region of linear extension as the Compton wavelength around the point of localization. In the massless limit ($M \rightarrow 0$), the wave function exhibits a power-law dependence $|\mathbf{r} - \mathbf{r}_0|^{-7/2}$ around the point of localization.

So far, we have ignored the vector character of the problem. For a transverse plasmariton we may try a three-vector momentum space wave function of the type

$$\psi_{\mathbf{r}_0}(\mathbf{p}; s) = [p^2 + (Mc_0)^2]^{1/4} \mathbf{e}_{\mathbf{p}s} \exp(-\frac{i}{\hbar} \mathbf{r}_0 \cdot \mathbf{p}), \quad (59)$$

where $\mathbf{e}_{\mathbf{p}s}$ ($s = 1$ or 2) is one of two transverse unit vectors. The longitudinal unit vector, which points in the \mathbf{p} -direction, is denoted by $\mathbf{e}_{\mathbf{p}3}$ below. When the gradient operator, $\nabla_{\mathbf{p}}$, acts on the wave function, it mixes in a longitudinal component, and this effect in itself makes it problematic (impossible) to take the $\hat{\mathbf{r}}$ -operator in Eq. (53) as the plasmariton position operator. Instead, it is perhaps natural to use the dyadic quantity

$$\hat{\mathbf{r}}_{ij} = i\hbar \left\{ \delta_{ij} \left(\nabla_{\mathbf{p}} - \frac{\mathbf{p}}{2[p^2 + (Mc_0)^2]} \right) - \sum_{s=1}^3 [\nabla_{\mathbf{p}}(\mathbf{e}_{\mathbf{p}s})_i] (\mathbf{e}_{\mathbf{p}s})_j \right\} \quad (60)$$

as a candidate for a plasmariton position operator.⁷ This choice leads to problems in configuration space, however, because the related wave function (Fourier integral transform of Eq. (59)) is not convergent. If one tries to use the standard regularization procedure the divergence-free demand on the wave function cannot be maintained.

The photon position operator problem cannot be detached from the spatial localization issue of photons emitted from a source (microscopic or mesoscopic). In the field-quantized description the relation between the transverse electric field operator, $\hat{\mathbf{E}}_T$, and the transverse current density operator, $\hat{\mathbf{J}}_T$, can be written as follows:

$$\hat{\mathbf{E}}_T(\mathbf{r}, t) = \mu_0 \int_{-\infty}^{\infty} g(|\mathbf{r} - \mathbf{r}'|, t - t') \cdot \frac{\partial}{\partial t'} \hat{\mathbf{J}}_T(\mathbf{r}', t') d^3 r' dt'. \quad (61)$$

If one now replaces $\hat{\mathbf{J}}_T$ by $\hat{\mathbf{J}}$ with the help of the transverse delta function the "propagator" relation

$$\hat{\mathbf{E}}_T(\mathbf{r}, t) = \mu_0 \int_{-\infty}^{\infty} \overleftrightarrow{\mathbf{D}}_T(\mathbf{r} - \mathbf{r}', t - t') \cdot \frac{\partial}{\partial t'} \hat{\mathbf{J}}(\mathbf{r}', t') d^3 r' dt' \quad (62)$$

emerges. Here, the transverse propagator has the explicit dyadic form⁸

$$\overleftrightarrow{\mathbf{D}}_T(\mathbf{R}, \tau) = -\frac{1}{4\pi R} \delta\left(\frac{R}{c_0} - \tau\right) (\overleftrightarrow{\mathbf{U}} - R^{-2} \mathbf{R}\mathbf{R}) + \frac{c_0^2 \tau}{4\pi R^3} \Theta(\tau) \Theta\left(\frac{R}{c_0} - \tau\right) (\overleftrightarrow{\mathbf{U}} - 3R^{-2} \mathbf{R}\mathbf{R}), \quad (63)$$

where Θ is the Heaviside step function. The form in Eq. (63) tells us that the transverse photon localization (in general) is limited to the extension of the rim zone, and that one cannot maintain the Einstein causality when compressing the source domain artificially from $\hat{\mathbf{J}}_T$ to $\hat{\mathbf{J}}$. Eq. (63) has served as a starting point for microscopic photon tunneling analyses in near-field electrodynamics.¹⁶

REFERENCES

- [1] Landau, L. and Peierls, R. *Zeitschr. Phys.* **62**, 188 (1930).
- [2] Oppenheimer, J. R. *Phys. Rev.* **38**, 725 (1931).
- [3] Keller, O. *Progr. Opt.* **50**, 51 (2007).
- [4] Bialynicki-Birula, I. *Progr. Opt.* **36**, 245 (1996).
- [5] Goldstein, H., [*Classical Mechanics*], Addison-Wesley, London (1980).
- [6] Born, M. and Wolf, E., [*Principles of Optics*], Cambridge Univ., Cambridge (1999).
- [7] Hawton, M. *Phys. Rev.* **A59**, 3223 (1999).
- [8] Keller, O. *Phys. Rep.* **411**, 1 (2005).

- [9] Keller, O. *Phys. Rev.* **A76**, 062110 (2007).
- [10] Keller, O. *Phys. Rev.* **A58**, 3407 (1998).
- [11] Keller, O. *Phys. Rev.* **A62**, 022111 (2000).
- [12] Keller, O. *Laser Part. Beams* **26**, 287 (2008).
- [13] Cohen-Tannoudji, C., Dupont-Roc, J., and Grynberg, G., [*Photons and Atoms. Introduction to Quantum Electrodynamics*], Wiley, New York (1989).
- [14] Mandl, F. and Shaw, G., [*Quantum Field Theory*], Wiley, Chichester (2001).
- [15] Weinberg, S., [*The Quantum Theory of Fields, Vol. I, Foundations*], Cambridge Univ., Cambridge (1996).
- [16] Keller, O. *Phys. Rev.* **A60**, 1652 (1999).